

ACCEPTED

FOR REGISTRATION

September 20, 2016

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

DOC ID: 546335



## SOFT BAIT

FOR INDOOR AND OUTDOOR USE

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT HOUSE MICE

KILLS OTHER RODENT SPECIES AS LISTED ON THIS LABEL

NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN ONE NIGHT'S FEEDING  
WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS

Active Ingredient: difethialone . . . . . 0.0025%

Inert Ingredients . . . . . 99.9975%

Total . . . . . 100.0000%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See inner panel for additional precautionary statements.

**LIPHA<sup>TM</sup> TECH**

Approx. 363 x 10g Pouches

This bait is packaged in 0.35 ounce  
(10 grams) individual packs

It is illegal to sell this product in individual containers holding less than 16 pounds of bait.

**Net Weight: 8 lbs. (3.6 kg) Pail**

UPC will be  
0 13549 52352 7

Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209  
800-351-1476

EPA Reg. No. 7173-258  
EPA Est. No. 7173-WI-1  
Product No. 82352  
Label No. 150-4979-0516

## INSTRUCCIONES DE USO

Es una contravención de la ley federal usar este producto de manera contraria a las instrucciones de la etiqueta.

**LEA ESTA ETIQUETA** y siga todas las instrucciones y precauciones. Utilícelo solamente para en los lugares, para las plagas y con los métodos de aplicación que se especifican en esta etiqueta.

**IMPORTANTE:** No exponga los raticidas donde haya niños, mascotas ni animales salvajes inofensivos que no son el objetivo.

Para ayudar a prevenir accidentes:

1. Almacene el producto que no esté en uso fuera del alcance de niños y mascotas.
2. Aplique el cebo en lugares fuera del alcance de niños, mascotas, animales domésticos y animales salvajes inofensivos que no son el objetivo, o bien póngalo en trampas de cebo resistentes a las alteraciones. Estas trampas deben ser resistentes a la destrucción por parte de perros y niños menores de seis años de edad y deben usarse de tal manera que eviten que dichos niños lleguen a los compartimentos del cebo y tomen contacto con él. Si el cebo puede salirse de las trampas al levantarlas, las unidades deben estar aseguradas o fijas de alguna otra manera. Se necesitan trampas todavía más resistentes en áreas abiertas a animales con pezuñas, mapaches, osos, otros animales potencialmente destructivos o en áreas propensas al vandalismo.
3. Descarte el recipiente del producto, producto restante, estropeado y cebo no consumido según se especifica en esta etiqueta.

**Nota:** Las estaciones de cebo son solamente para uso en exteriores, sobre la superficie. Se deben utilizar estaciones de cebo resistentes a las alteraciones si hay niños, mascotas, mamíferos que no quiera perjudicar o aves que podrían tener acceso al cebo.

**RESTRICCIONES DE USO:** Este producto puede usarse solo para controlar ratas de Noruega, ratas de tejado, ratones domésticos, Ratón ciervo (*Peromyscus maniculatus*), ratón de patas blancas (*Peromyscus leucopus*), Ratón de algodónero (*Peromyscus gossypinus*), ratá algodonera (*Sigmodon hispidus*), ratón cosechero del este\* (*Reithrodontomys humilis*), ratón dorado\* (*Ochrotomys nuttalli*), rata polinesia\* (*Rattus exulans*), campañol del prado\* (*Micromys pennsylvanicus*), rata del cuello blanco\* (*Neotoma albigena*), rata del llano sureño\* (*Neotoma micropus*) y rata mexicana\* (*Neotoma mexicana*) en y dentro de un perímetro de 100 pies de estructuras erigidas por el hombre y construidas de manera tal que sean vulnerables a invasiones de roedores comensales, o bien, para rodear o atacar plagas de roedores. Entre dichas estructuras se encuentran las casas y otras residencias permanentes o temporales, centros procesadores de alimentos, edificios industriales y comerciales, recipientes de basura, edificios agrícolas y públicos, vehículos de transporte (buques, trenes, aeronaves), muelles y edificios portuarios o terminales, así como estructuras relacionadas alrededor de estos lugares y asociadas con ellos. Se prohíbe colocar cebos y cercas en perímetros que estén a más de 100 pies de una estructura, según se define arriba. No venda este producto en envases individuales con menos de 16 libras de cebo. No lo coloque cerca o dentro de las aberturas de conductos de ventilación. No contamine el agua, la comida, el alimento para animales o los equipos de manejo de la alimentación, o la leche o los equipos para manipular carne. No lo aplique directamente a alimentos o forrajes. No esparza el cebo. Está prohibida la colocación de cebos de este producto en madrigueras.

\*No permitido para el uso contra las siguientes especies en California: rata algodonera, ratón cosechero del este, ratón dorado, rata polinesia, campañol del prado, rata de cuello blanco, rata del llano sureño, y rata mexicana.

**Selección de las áreas de tratamiento:** Determine las áreas donde es más probable que encuentren y consuman el cebo las ratas o ratones. En general, es a lo largo de paredes, junto a aberturas mordidas, en rincones y lugares ocultos, entre pisos o paredes, al lado de madrigueras, entre pisos o paredes, o en ubicaciones donde se hayan visto roedores o señales de su presencia. Retire todos los alimentos alternativos que sea posible. (041216)

## DECLARACIONES PREVENTIVAS

### Peligroso para personas y animales domésticos

**PRECAUCIÓN:** es perjudicial si se traga o absorbe a través de la piel. Causa irritación moderada en los ojos. Evite el contacto con la piel, los ojos o la ropa. Manténgalo alejado de los niños, animales domésticos y mascotas. Todos aquellos que manipulen el producto (incluidos quienes lo apliquen) deben usar pantalones largos, zapatos, calcetines, y guantes impermeable. Toda persona que reciba animales muertos o cebos sin utilizar después de la aplicación de este producto debe usar guantes impermeables.

**Requisitos Para la Seguridad del Usuario:** Siga las instrucciones del fabricante para limpiar y mantener el equipo de protección personal (EPP). Si no hay instrucciones para las prendas lavables, utilice detergente y agua caliente. Mantenga y lave el EPP aparte del resto de la ropa para lavar. Quitese el EPP inmediatamente después de manipular este producto. Lave la parte exterior de los guantes antes de quitárselos. Tan pronto como pueda, lávese las manos minuciosamente después de la aplicación de los cebos y antes de comer, beber, mascar goma, usar tabaco o usar el baño, y póngase ropa limpia.

### PRIMEROS AUXILIOS

Tenga esta etiqueta a mano cuando le aconsejen un tratamiento.

**Si se ingiere:** Llame a un centro de control de envenenamiento o a un doctor de manera inmediata para que le aconsejen un tratamiento. Si la persona puede tragar, haga que beba un vaso de agua. No induzca el vómito a menos que se lo indique el centro de control de envenenamiento o el médico.

**Si hay contacto con los ojos:** Mantenga abierto el ojo y enjuague lenta y suavemente con agua durante 15 a 20 minutos. Quitese los lentes de contacto, en caso de usarlos, después de los primeros cinco minutos y luego continúe enjuagando. Llame a un centro de control de envenenamiento o a un doctor para que le aconsejen un tratamiento.

**Si entra en contacto con la piel o la ropa:** Quitese la ropa contaminada. Enjuague la piel con abundante agua fría durante 15 a 20 minutos. Llame a un centro de control de envenenamiento o a un doctor para que le aconsejen un tratamiento.

**TRATAMIENTO POR ENVENENAMIENTO DE MASCOTAS:** Si el animal come algún cebo, llame al veterinario de inmediato.

**NOTA PARA EL MÉDICO O VETERINARIO:** Contiene el anticoagulante difethylone. En caso de ingestión, este material puede reducir la capacidad de coagulación de la sangre y producir una hemorragia. En el caso de seres humanos o animales que ingieran cebos o muestran signos de envenenamiento (hemorragias o tiempos prolongados de protrombina), suministrelles vitamina K1 por inyección intramuscular o en forma oral. También en el caso de las mascotas, si es necesario, verifique el tiempo de protrombina cada tres días hasta que los valores vuelvan a la normalidad (hasta 30 días). En casos graves, se deberán realizar transfusiones de sangre.

**RIESGOS AMBIENTALES:** Este producto es extremadamente tóxico a los mamíferos y pájaros. Los perros, gatos y otros mamíferos y pájaros predadores o carroñeros pueden sufrir envenenamiento si se alimentan de animales que hayan comido el cebo. No lo aplique directamente al agua o en áreas donde haya agua superficial, o en zonas intermareales que estén debajo de la marca de agua máxima promedio. No contamine el agua por medio del lavado del equipo o el desecho de residuos.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL** and follow all use directions and precautions. Use only for the sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides.

To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats, house mice, Deer mouse (*Peromyscus maniculatus*), White-footed mouse (*Peromyscus leucopus*), Cotton mouse (*Peromyscus gossypinus*), Cotton rat\* (*Sigmodon hispidus*), Eastern harvest mouse\* (*Reithrodontomys humulis*), Golden mouse\* (*Ochrotomys nuttalli*), Polynesian rat\* (*Rattus exulans*), Meadow vole\* (*Microtus pennsylvanicus*), White-throated woodrat\* (*Neotoma albigena*), Southern plains woodrat\* (*Neotoma micropus*) and Mexican woodrat\* (*Neotoma mexicana*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings, and related structures around and associated with these sites. Fence and perimeter baiting beyond 100 feet from a structure, as defined above, is prohibited. Do not sell this product in individual containers holding less than 16 pounds of bait. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not apply directly to food or feed crops. Do not broadcast bait. Burrow baiting with this product is prohibited.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat.

**Selection of Treatment Areas:** Determine areas where rats and/or house mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls, beside burrows, or in locations where rats and/or house mice or their signs have been observed. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each pouch contains 0.35 oz. (10g). **House Mice:** Apply 1 pouch per placement, usually spaced 8 to 12 feet apart. Up to 4 pouches may be needed at points of very high house mouse activity. Maintain a constant supply of fresh bait for 15 days or until signs of house mouse activity cease. **Rats:** Apply 9 to 43 pouches per placement, usually spaced 15 to 30 feet apart. Maintain a constant supply of fresh bait for 10 days or until signs of rat activity cease. Bait may be removed from pouch and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash. **Follow-up:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

(041216)

## PRECAUTIONARY STATEMENTS

### Hazard to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear long pants, shoes, socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet, and change into clean clothing.

### FIRST AID

Have this label with you when obtaining treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor immediately for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin with plenty of cool water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.

### TREATMENT FOR PET POISONING:

If animal eats bait, call veterinarian at once.

**NOTE TO PHYSICIAN OR VETERINARIAN:** Contains the anticoagulant difethialone. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. For humans or animals ingesting bait and/or showing obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K<sub>1</sub> intramuscularly or orally. Also for pets, if needed, check prothrombin time every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of in trash or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty container, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(CONTINUADO)

## INSTRUCCIONES DE APLICACIÓN

**Cada paquete contiene 0.35 onzas (10g).** **Ratones domésticos:** Aplique un paquete por punto de colocación, espaciaron generalmente 8 a 12 pies (2.5 a 3.7 metros) aparte. Puede que se necesiten hasta cuatro paquetes en puntos donde haya una actividad muy alta de ratones. Mantenga un suministro constante de cebo fresco durante 15 días o hasta que cesen las señales de actividad de ratones domésticos.

**Ratas:** Aplique entre 9 a 43 paquetes por punto de colocación, espaciaron generalmente 15 a 30 pies (4.5 a 9 metros) aparte. Mantenga un suministro constante de cebo fresco durante 10 días o hasta que cesen las señales de actividad de ratas. El cebo se puede quitar de bolsa y colocar en un dispositivo de alimentación que asegure el cebo desempaquetado en una estación del cebo. Deseche la envoltura vacía en la basura.

**Seguimiento:** Reemplace el cebo contaminado o descompuesto inmediatamente. Utilice guantes impermeables para recolectar y eliminar todos los animales muertos visibles y el cebo restante, como también cualquier cebo que pueda quedar expuesto ante aquellos animales que no quiera perjudicar. Para evitar la reinfestación, limite lo más posible la disponibilidad de agua, lugares aptos para madrigueras y alimentos que atraigan ratones. Si ocurre la reinfestación, repita el tratamiento. Para una infestación continua, instale trampas de cebo permanentes y reabastezca el cebo según se necesite. (041216)

## ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, la comida o el forraje mediante el almacenamiento o la eliminación.

**Almacenamiento del pesticida:** guárdelo en el envase original, en un lugar fresco y seco fuera del alcance de los niños y las mascotas.

**Eliminación del pesticida:** los desechos que se generen por el uso de este producto pueden eliminarse en la basura o en una instalación aprobada para la eliminación de desechos.

**Manipulación del envase:** envase no recargable. No vuelva a utilizarlo ni a recargarlo. Cuando el envase esté totalmente vacío, ofrézcalo para reciclarlo o reacondicionamiento; o bien, píñchelo y deséchelo en un vertedero sanitario.

**GARANTÍA:** En la medida que cumpla con la ley aplicable, el vendedor no concede garantías, expresas ni implícitas, referentes al uso de este producto aparte de lo indicado en la etiqueta. El comprador asume todos los riesgos del uso y/o la manipulación de este material cuando dicho uso y/o manipulación sea contraria a las instrucciones de la etiqueta.

6.875

Printed Color



## SOFT BAIT

### FOR INDOOR AND OUTDOOR USE

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE  
AND WARFARIN-RESISTANT HOUSE MICE

KILLS OTHER RODENT SPECIES AS LISTED ON THIS LABEL

NORWAY RATS AND HOUSE MICE MAY CONSUME  
A LETHAL DOSE IN ONE NIGHT'S FEEDING  
WITH FIRST DEAD RODENTS APPEARING  
FOUR OR FIVE DAYS AFTER FEEDING BEGINS

Active Ingredient: difethialone .....	0.0025%
Inert Ingredients .....	99.9975%
Total .....	100.0000%

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

See side panel for additional precautionary statements.

Please refer to booklet label for complete directions for use.  
This section is NOT a complete product label. If the label booklet has been removed from this container, contact Liphatech, Inc. for a replacement.

Approx. 363 x 10g Pouches

This bait is packaged in 0.35 ounce  
(10 grams) individual packs

It is illegal to sell this product in individual containers holding less than 16 pounds of bait.

**LIPHA TECH**®  
Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209  
(800) 351-1476

EPA Reg. No. 7173-258

Product No. 82352

EPA Est. No. 7173-WI-1

Label No. 150-4979-0516

### PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

## CAUTION:

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear long pants, shoes, socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet, and change into clean clothing.

**FIRST AID:** Have this label with you when obtaining treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor immediately for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin with plenty of cool water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.

**TREATMENT FOR PET POISONING:** If animal eats bait, call veterinarian at once.

**NOTE TO PHYSICIAN OR VETERINARIAN:** Contains the anticoagulant difethialone. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. For humans or animals ingesting bait and/or showing obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K<sub>1</sub> intramuscularly or orally. Also for pets, if needed, check prothrombin time every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of in trash or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty container, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill.

**Net Weight: 8 lbs. (3.6 kg) Pail**

5.0"

Printed Color

## PRECAUTIONARY STATEMENTS

### Hazard to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear long pants, shoes, socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

**User Safety Requirements:** Follow manufacturer's instructions for clearing/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet, and change into clean clothing.

### FIRST AID

Have this label with you when obtaining treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor immediately for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin with plenty of cool water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.

**TREATMENT FOR PET POISONING:** If animal eats bait, call veterinarian at once.

**NOTE TO PHYSICIAN OR VETERINARIAN:** Contains the anticoagulant difethialone. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. For humans or animals ingesting bait and/or showing obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K<sub>1</sub> intramuscularly or orally. Also for pets, if needed, check prothrombin time every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. (041216)

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of in trash or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty container, then dispose of empty container in trash or at an approved waste disposal facility.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



### FOR INDOOR AND OUTDOOR USE

### KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT HOUSE MICE

### KILLS OTHER RODENT SPECIES AS LISTED ON THIS LABEL

### NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN ONE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS

Active Ingredient: difethialone .....	0.0025%
Inert Ingredients .....	99.9975%
Total .....	100.0000%

### KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

**LIPHA TECH®**

Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209  
800-351-1476

**It is illegal to sell this product in individual containers holding less than 16 pounds of bait.**

### Approx. 726 x 10g Pouches

This bait is packaged in 0.35 ounce (10 grams) individual packs

**Net Weight: 16 lbs. (7.3 kg) (8 lbs. x 2) Case**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ THIS LABEL and follow all use directions and precautions. Use only for the sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats, house mice, Deer mouse (*Peromyscus maniculatus*), White-footed mouse (*Peromyscus leucopus*), Cotton mouse (*Peromyscus gossypinus*), Cotton rat\* (*Sigmodon hispidus*), Eastern harvest mouse\* (*Reithrodontomys humulis*), Golden mouse\* (*Ochrotomys nuttalli*), Polynesian rat\* (*Rattus exulans*), Meadow vole\* (*Microtus pennsylvanicus*), White-throated woodrat\* (*Neotoma albigena*), Southern plains woodrat\* (*Neotoma micropus*) and Mexican woodrat\* (*Neotoma mexicana*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings, and related structures around and associated with these sites. Fence and perimeter baiting beyond 100 feet from a structure, as defined above, is prohibited. Do not sell this product in individual containers holding less than 16 pounds of bait. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not apply directly to food or feed crops. Do not broadcast bait. Burrow baiting with this product is prohibited.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat.

**Selection of Treatment Areas:** Determine areas where rats and/or house mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls, beside burrows, or in locations where rats and/or house mice or their signs have been observed. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each pouch contains 0.35 oz. (10g). **House Mice:** Apply 1 pouch per placement, usually spaced 8 to 12 feet apart. Up to 4 pouches may be needed at points of very high house mouse activity. Maintain a constant supply of fresh bait for 15 days or until signs of house mouse activity cease.

**Rats:** Apply 9 to 43 pouches per placement, usually spaced 15 to 30 feet apart. Maintain a constant supply of fresh bait for 10 days or until signs of rat activity cease. Bait may be removed from pouch and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash.

**Follow-up:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinestation, limit sources of rodent food, water and harborage as much as possible. If reinestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

EPA Reg. No. 7173-258


EPA Est. No. 7173-WI-1

Product No. 8232

Label No. 150-4974-0516



0 13549 82352 8



Live  
Print  
Area  
 $W=1.75 \times H=1.375$

